

# New horizons in platelet research: Understanding and harnessing platelet functional diversity

Ilya Reviakine\*

*Institute of Functional Interfaces (IFG), Karlsruhe Institute of Technology (KIT), Hermann-von Helmholtz-Platz, Eggenstein-Leopoldshafen, Germany*

**Abstract.** Recent years have been ripe with discoveries of non-haemostatic platelet functions. This led to the appreciation of the significant, previously unknown, role played by the platelets in various pathologies and regenerative processes. As a result, exciting opportunities for clinical applications in fields as diverse as regenerative medicine and cancer treatment are emerging. However, their realization depends on the understanding of the regulatory mechanisms governing these diverse platelet functions, so that particular platelet responses could be artificially tailored to specific clinical situations. Current understanding of the signalling pathways controlling haemostatic responses is rooted in the development of quantitative assays for measuring them and sensitive markers for their quantification. However, the existing assays and markers are not sufficiently sensitive for distinguishing between individual signalling pathways and unravelling inter-pathway connections. Moreover, entirely new approaches are needed for studying non-haemostatic platelet functions, since there are currently no assays or markers for quantifying them. We review the on-going efforts in these directions, including our own recent work on using lectins as sensitive probes for profiling platelet activation.

## 1. Platelets and their therapeutic potential

Last decade in platelet research has been ripe with exciting surprises. Originally thought of as clotting factories (Fig. 1), platelets turned out to be involved in a wide variety of physiological and pathophysiological processes [75]. They appear to function as a Swiss army knife, unsheathed as a first-response to injury, in processes ranging from coagulation, where they limit traumatic blood loss, to inflammation and the subsequent steps of the wound healing cascade [5, 88, 97], angiogenesis [88, 96], vascular remodelling and maintenance of vascular integrity [55], *de novo* blood vessel synthesis, initial response to pathogens [148], and other aspects of both innate and adaptive immune response (Fig. 2) [88, 121, 138]. Misfiring of these mechanisms is implicated in pathological conditions as diverse as atherosclerosis [13, 67, 78, 110] and the development of cancer metastases [33, 34, 127]. This diversity raises important questions but also presents therapeutic opportunities that are as unique as they are varied. The realization of these opportunities is predicated on the understanding of the regulatory processes that govern platelets' various functional roles.

---

\*Corresponding author: Ilya Reviakine, Institute of Functional Interfaces (IFG), Karlsruhe Institute of Technology (KIT), Hermann-von Helmholtz-Platz 1, 76344 Eggenstein-Leopoldshafen, Germany. E-mail: [ilya.reviakine@kit.edu](mailto:ilya.reviakine@kit.edu).




Fig. 1. Platelets in Haemostasis. Under normal physiological conditions, platelets (large grey circles) circulate in the blood in a resting state. They do not aggregate and their interactions with the healthy blood vessel endothelium or normally folded plasma proteins are limited. Exposure of collagen and collagen-adsorbed von Willebrand factor at wound sites (a star-shaped, black defect in the grey vessel wall) leads to platelet adhesion. This is followed by activation that involves a conformational change of GPIIb/IIIa, a constitutively expressed integrin receptor on the platelet surface (small green circles on the platelet surface). Activated GPIIb/IIIa (green ellipses on the platelet surface) becomes competent to bind fibrinogen. This leads to platelet aggregation and plug formation. Concurrent secretion of granule contents (dashed white arrows), that includes platelet agonists such as ADP, and production of thromboxane, causes activation and recruitment of additional platelets to the growing platelet plug. Phosphatidyl serine (PS, red arrowheads) that becomes exposed on the surfaces of the activated platelets catalyses the assembly of the clotting factor complexes such as tenase and prothrombinase (not shown). Their activity leads to thrombin production and ultimately fibrin formation from fibrinogen. Fibrin polymerizes forming the clot that seals the wound. It also becomes a natural scaffold for the subsequent wound healing process, in which platelets also play a leading role.

From a historical perspective, it is interesting to note that evidence of platelet functional diversity dates back to some very early studies. For example, their role in maintaining vascular integrity was noted as early as 1940s by Danielli [21], and their involvement in the early stages of atherosclerosis [67, 78] is somehow connected to this role. In 1960s, the effect of platelets on vascular integrity was essentially re-discovered and investigated more systematically (see, e.g., Gimbrone et al. [36]; similarly, in his review, written in 1964, T.H. Spaet refers to this platelet effect as “well-recognized”, without, however, mentioning Danielli’s work [125]). Subsequently, some of the underlying mechanisms began to be clarified. Most recent ideas on the subject are summarized in refs. [55] and [139]. Both direct interactions between the platelets and the endothelial cells (adhesion) and indirect effects mediated by the platelet secretion reactions appear to be needed for maintaining the integrity of the inflamed vascular endothelium [55], where platelet relevance has been demonstrated in thrombocytopenic mice [37]. Quiescent platelets were shown to adhere and roll on activated endothelium by intravital microscopy [29, 139]. This process is mediated by the endothelial P-selectin and vWF, stored in the Weibel-Palade bodies [141] of the endothelial cells, that interact with constitutively expressed P-selectin glycoprotein ligand (PGSL-1) and GPIb, respectively [139]. Until now, it appears that some form of activation and expression of P-selectin on platelets is required for attracting leukocytes, however [139]. Key signalling events that mediate platelet adhesion to the inflamed endothelium, the details of adhering platelet activation, and the transformation to the putative state in which adhering platelets promote leukocyte extravasation




Fig. 2. Multifunctional platelets. Platelets appear to function as a Swiss army knife of first responses to injury, being involved in a variety of defensive reactions such as coagulation and subsequent wound healing processes, pathogen recognition and other aspects of the adaptive immune response, angiogenesis and vascular remodelling, and maintenance of vascular integrity. Pathologies such as thrombosis, atherosclerosis (not shown), and cancer metastases arise as a result of the malfunction of these responses. To explain these diverse functions of the platelets, it is supposed that platelets exhibit multiple activation states (indicated with question marks), as opposed to a unique procoagulant one (indicated on the left). Characteristics of these states are not yet understood. Colours and symbols are the same as in Fig. 1.

without initiating thrombus formation, remain obscure. This last process is thought to provide the link to the aetiology of atherosclerosis [67, 78]. *In vitro* studies of these processes continue to remain challenging [73].

Similarly, the link between platelets and cancer can be traced back to the work of Trousseau in 1860s; a very informative historical review of Trousseau's work can be found in ref. [70], while the molecular details of platelet-cancer links have recently been reviewed by Gay and Felding-Habermann [34]. Recent studies point to the relevance of their ability to assist extravasation of cells, their role in angiogenesis, and vascular remodelling. These platelet effects may or may not be directly linked to their procoagulatory function.

It is becoming clear that the procoagulant role of platelets may be divorced from their other functional roles, such as that of maintaining vascular integrity, angiogenesis, and vascular remodelling. All of this implies a multiplicity of different activation states. How these activities are regulated is not understood. Key questions surround relevant molecular triggers and specific responses, such as secretion and expression of specific signalling molecules, involved in the various roles platelets play. Understanding these mechanisms and harnessing them to stimulate the regenerative functions of the platelets while minimizing thrombotic and detrimental inflammatory effects would revolutionize prevention and treatment of cardiovascular disorders, the underlying atherosclerosis, healing of recalcitrant wounds, integration of implants, and possibly, even the treatment of cancers. The idea here is to tailor-manage platelet activity for different pathologies in individual patients. Such an approach to platelet-based therapy is quite different from the strategies for modulating platelet procoagulant activity constituting modern antiplatelet therapies [8, 60].

Current therapeutic approaches focus on taming the unwanted procoagulant activity of platelets. They entail targeting platelet surface receptors or modulating their autocrine loops. Aspirin remains the most

wide-spread and successful antiplatelet drug for chronic and prophylactic use [91, 100]. It is a cyclooxygenase inhibitor that blocks arachidonic acid metabolism into thromboxane A<sub>2</sub> (TXA<sub>2</sub>). TXA<sub>2</sub> is an important platelet agonist, and, as discussed below, a key element of platelet autoactivation. The second most successful class of antiplatelet agents are ADP receptor antagonists ticlopidine and clopidogrel, prasugrel, and ticagrelor. Introduction of the dual aspirin/ticlopidine (now replaced by clopidogrel) antiplatelet therapy in the late 90s has, without a doubt, revolutionized interventional cardiology by allowing the wide-spread use of stenting for treating coronary artery occlusion [16, 117]. Schömig et al. [118] and Holmes et al. [54] further discuss the combination of the dual antiplatelet therapy with oral anticoagulation, recommended, in particular, for patients with mechanical heart valves and other similar devices (e.g., LVADs), and a number of other conditions that lead to aggressive thrombosis, such as atrial fibrillation.

The efficiency of antiplatelet agents in modulating platelet activity is limited by the bleeding risk. For this reason, the development of oral GPIIb/IIIa antagonists met with some difficulties: they caused serious bleeding complications [60, 91]. This can be rationalized by the fact that GPIIb/IIIa represents the key point of intersection of many of the platelet signalling pathways (discussed in more detail below), is unique in promoting platelet aggregation by fibrinogen, and stabilizing clot formation. Its inhibition, therefore, works in a nearly all-or-none fashion, limiting the so-called therapeutic window of the drugs (the range of doses where platelet function modulation is maximized while the bleeding risk is minimized). GPIIb/IIIa antagonists are used in acute situations (e.g., during percutaneous coronary intervention), delivered by intravenous or intracoronary injections [42]. These, and other antiplatelet agents, together with the promising developments in this area, have been recently comprehensively reviewed by Jackson and Schoenwaelder [60], K. Schrör [119], Bhatt and Topol [8], and Papp et al. [91]. The issues of resistance to aspirin and some of the other antiplatelet therapies are discussed by Fitzgerald and Pirmohamed [27], M. Cattaneo [11], and Papp et al. [91].

When discussing antiplatelet therapeutic agents, the topic of inflammation cannot be avoided. Firstly, platelets are themselves inflammatory mediators [8, 39, 88, 121, 138]. Secondly, aspirin is well-known for its anti-inflammatory and analgesic effects—in fact, originally this was its primary therapeutic use (see ref. [147] for a recent review). This has to do, of course, with its effect on both of the cyclooxygenase isoforms, COX-1 and COX-2. Its antiplatelet effects (mediated by COX-1 inhibition) are apparent at lower doses than the anti-inflammatory ones (mediated primarily by COX-2 inhibition) [91]. Similarly, GPIIb/IIIa antagonists such as Abciximab exhibit anti-inflammatory effects through cross-reactivity with integrin receptors expressed on macrophages [79]. Other antiplatelet agents, such as clopidogrel, appear to exhibit some anti-inflammatory properties, mediated by their inhibition of platelet activity—but this aspect is much less well understood [32].

Cardiovascular diseases (CAD) are currently responsible for more deaths than any other disease [123, 129]. Critical role of platelets in CAD is underscored by the success of antiplatelet therapies in mitigating the consequences of CAD, minimizing the risk of serious complications and death, and of the effects of the disease on patients' lifestyles [60, 91, 100, 123]. The severity of the CAD epidemic dictates the need for the development of new and improved antiplatelet agents [60, 123]. Current strategies, however, continue to focus on the haemostatic functions of platelets. Approaches harnessing their regenerative potential are yet to come.

Efforts to harness the regenerative potential of platelets are already under way in other areas. On one hand, Davies and colleagues observed a correlation between the extent, to which dental implant materials activated platelets, and osteoconductivity displayed by the implants [68, 92, 93]. On the other hand, spurred on by the surgical applications of fibrin gels as surgical glues [61], autologous

platelet-containing fibrin gels started to be used in dental, oral, maxillofacial, and plastic surgery [4, 10, 24, 88, 145]. Reports of their successful applications, however, remain somewhat anecdotal. The key hypothesis that appears to be driving these efforts is that growth factor secretion by the platelets enhances wound healing [6]. This hypothesis is difficult to reconcile with the fact that platelets secrete growth factors with contradictory functions (see below). Mechanistic studies are sparse and numerous open questions remain. In particular, the absence of leukocytes from the platelet-rich plasmas used in therapeutic applications may have an effect on the wound healing process initiated by these products. These doubts notwithstanding, the application of platelet-rich plasmas to wound healing and implant integration is an interesting example of a therapeutic application of platelets.

## 2. Platelets and their haemostatic activity

Under normal physiological conditions, platelets circulate in a quiescent state. The ability of quiescent platelets to interact with each other, with the normally folded plasma proteins, and with other cells, is limited (Fig. 1). On the contrary, activated, procoagulant platelets aggregate, adhere to other cells, and interact with plasma proteins such as fibrinogen and the factors of the coagulation cascade.

Their ability to aggregate arises from a change in the conformation of the GPIIb/IIIa integrin complex, which then becomes competent to bind fibrinogen that cross-links multiple platelets. As discussed above, this made GPIIb/IIIa an obvious target for antiplatelet medication, and several drugs targeting this receptor were developed. As already mentioned, they are used intravenously or intracoronary in acute situations, but carry with them too high a bleeding risk for prolonged use in chronic situations.


As a part of the activation process, platelets also increase in size, exposing the area previously stored in membrane folds of the open canalicular system (OCS), and change shape. Phospholipid phosphatidyl serine (PS), that activated platelets express on their surface, serves as an anchor for the assembly of clotting factor complexes (tenase and prothrombinase) that ultimately catalyse thrombin formation [7, 35, 85, 135, 151]. Thrombin cleaves fibrinogen into fibrin, and the latter polymerizes to form a clot. As PS expression is also associated with apoptosis, understanding the role apoptotic pathways play in platelet activation became an interesting avenue of investigation [46, 76]. Last but not least, activated platelets release the content of their secretory granules. Dense granules contain small molecules (ADP, serotonin, calcium, etc.) while alpha granules contain proteins, more than 300 growth factors, cytokines, adhesion mediators, and clotting factors [88, 95, 146, 150].

Details of these processes have been studied intently since the 1960s, in what B. S. Coller in his review [16] refers to as the “mechanistic period” in platelet research. The impetus for these studies was given by the development of quantitative assays for measuring platelet activity in coagulation and the discovery of reliable platelet activation markers. These included the platelet adhesion assay, aggregometry, aggregometry coupled with the luciferase assay to quasi-simultaneously measure granule secretion and aggregation, and subsequently—flow cytometry using monoclonal antibodies against platelet surface activation markers. These were later combined with intracellular calcium measurements [16]. Notwithstanding the impact of these tools on unravelling the molecular building blocks underlying platelet physiology, they do have limitations. Firstly, it appears that they are not sufficiently sensitive to distinguish between individual signalling pathways. Secondly, they are not sensitive to non-haemostatic platelet functions. In other words, both unravelling individual signalling pathways in platelets and understanding other aspects of platelet function will require the development of a new “toolbox” of assays and markers.

### 3. Platelet signalling pathways

The signalling circuitry that controls platelet activation processes is impressively complex and not completely understood. A separate review would be required to cover this topic, and indeed, several are available [16, 17, 22, 60, 77, 80, 90, 136, 137]. Information is also available in textbooks [84]. We will limit ourselves to a very much simplified sketch (Fig. 3) that serves to illustrate some key features.

The extent, to which platelet function is fine-tuned, can be appreciated from the number of different agonists and modulators that control it. These include molecules expressed on subendothelial tissues (collagen, collagen-adsorbed vWF, fibrin), and soluble substances that are secreted or produced by other cells (ADP [30], epinephrine, neuromediators [114], platelet activating factor (PAF) [23, 52]), the platelets themselves (serotonin, ADP, ATP, thromboxane), generated as a part of the coagulation cascade (thrombin, the most potent of platelet agonists), or constitutively present in the plasma (plasma vWF becomes an agonist under conditions of high shear [116]; the role of vWF in platelet aggregation and adhesion at different shear rates was described by Weiss et al. as far as 1978 in what appears to be one of the first reports on the subject [142]. See also ref. [58] for an interesting review). For several of the agonists, there is more than one receptor: P2Y1 and P2Y12 for ADP; [62, 90] protease activated receptors PAR1 and PAR4, [17] as well as GPIb, [109] for thrombin [22]. Some of these are shown in Fig. 3. These receptors have different binding affinities for their agonists and contribute to platelet responses in different ways [3, 18, 62, 128]. Current list of platelet agonists, and their receptors, is most likely incomplete—a point aptly illustrated by the recent account of the role galectins play in platelet activation [113].



Downstream of the soluble agonist receptors are G-proteins: Gq, Gi, and G12/13 (reviewed in refs. [14, 80, 89]). Of note, PAR1 receptor for the strongest platelet agonist thrombin, is connected to all three G-proteins, while the two ADP receptors are connected to one kind of G-proteins each [89]. Collagen and vWF receptors are kinases and act through the tyrosine kinase signalling pathways (Fig. 3) [77].

A key element of platelet signalling pathways is the intracellular calcium rise that follows the interaction of most of the agonists with their receptors. Calcium rise is initiated by the phospholipase C (PLC) hydrolysis of phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>) that generates IP<sub>3</sub>. IP<sub>3</sub> induces Ca release from intracellular stores into the cytoplasm. The depletion of calcium in the stores triggers its entry from the extracellular space through the STIM1 (stromal interaction molecule 1)/Orai1 Ca-sensing and transport system. This process, called store-operated calcium entry (SOCE, Fig. 3), generates a rise in the intracellular calcium from nM range to  $\mu$ M range; it is described in detail in the reviews by Varga-Szabo et al. [136, 137] There are several forms of PLC; in particular, PLC $\beta$  is activated by a G-protein (Gq), while PLC $\gamma$ 2 is activated by the tyrosine kinase pathway (Syk) downstream of collagen and vWF receptors [136, 137].

It should be noted that platelet activation can proceed in the absence of extracellular calcium—as studies performed in the presence of citrate or EDTA indicate [38, 47, 57, 115]. EDTA is known to

---

Fig. 3. Some aspects of platelet regulatory pathways are illustrated in this figure. Examples of soluble agonists shown are thrombin, ADP, and thromboxane. Their receptors are coupled to G proteins (GPCRs), the identity of which is indicated below each receptor. Agonists that are typically exposed at surfaces of cells or tissues are represented by vWF and collagen. Their receptors act through the tyrosine kinase pathways (e.g., Syk) [51, 69, 80]. (Collagen also binds to GPIa/IIa, also known as integrin  $\alpha_{IIb}\beta_1$ , and CD36; these are not shown). The fibrinogen receptor, GPIIb/IIIa ( $\alpha_{IIb}\beta_{III}$  integrin), is a key point of intersection of all of the platelet regulatory pathways involved in procoagulant response (tyrosine kinase, G-protein mediated, etc.) This protein controls the aggregation of platelets into the platelet plug and the thrombus. A prominent second messenger system, Store-Operated Calcium Entry (SOCE), is indicated as a blue hexagon. It is reasonably well-understood and described in detail in Varga-Sabo et al. [136, 137] SOCE is accessed through PLC isoforms; PLC $\beta$ , activated by the G-protein Gq, and PLC $\gamma$ , that is activated via the Syk pathway [136, 137]. Blue arrowheads follow the ADP-mediated pathway from its interaction with the P2Y<sub>1</sub> and P2Y<sub>12</sub> receptors to the Ca-mediated shape change, GPIIb-IIIa activation, aggregation- and Ca-dependent PLA<sub>2</sub> activation and TXA<sub>2</sub> production. It should be noted that ADP activates integrin  $\alpha_{IIb}\beta_1$ , making it competent for high-affinity collagen binding, in the same way it activates GPIIb-IIIa for fibrinogen binding [64]. Purple arrows follow the TXA<sub>2</sub> pathways that lead to granule secretion (potentiated by ADP through the P2Y<sub>12</sub> receptor). Some of the secretion events are Ca-dependent, while others are not. Black arrow illustrates the effect vWF binding to the GPIb-IX-V complex has on the secretion of ADP [83]. Full brown arrow indicates the calcium-independent pathway connecting GPVI with the granule secretion, and TXA<sub>2</sub> production. It is modulated by ADP through its action on the P2Y<sub>12</sub> receptor that is coupled to the Gi protein. Dashed brown arrow indicates a pathway terminating in the cytoplasmic PLA<sub>2</sub> (cPLA<sub>2</sub>), another Ca-dependent phospholipase. Both vWF/GPIb-IX-V and collagen/GPVI directly activate GPIIb/IIIa. Grey arrow indicates the Syk activation of PLC $\gamma$ 2, which may be responsible for collagen-induced calcium rise in aspirin-treated platelets. A dotted brown arrow indicates collagen-induced, calcium-dependent loss in the PS asymmetry. This appears to be a synergistic pathway dependent on both the particular shape of the calcium rise signal and the activation of the tyrosine kinase pathway [51]. Isoforms of protein kinase C (PKC), activated by DAG and Ca, are involved in the regulation of the secretion events, independently of the pathway [49]. Gi proteins activate PI3K. G12/13 act through the rho proteins (not shown). Thrombin is the most potent of the platelet agonists, and achieves most of the effects of the other agonists—shape change, calcium rise, GPIIb/IIIa activation and aggregation, and secretion. Its PAR1 receptor is coupled to all three of the G-proteins, PAR4—to two of the three. In comparison, ADP receptors are coupled to one each, and TXA<sub>2</sub> receptor is coupled to two. Furthermore, some of the effects of thrombin on the platelets are also exerted through GPIb, thus covering the entire set of the platelet activation pathways. This partially explains why thrombin covers most of the range of the responses induced by other agonists. The individual arrowheads corresponding to thrombin signals are not shown, as they would considerably complicate the diagram. The details of many of the pathways are still being worked out. This figure was adapted from ref. [43].



damage platelets; White et al. investigated which of its effects were reversible and which were not [143]. Similarly, extracellular calcium is not a platelet agonist; this appears to be a source of some confusion, especially in the biomaterials field, apparently because coagulation may be induced by recalcification of citrated blood or platelet-rich plasma. The point here is that platelet aggregation and coagulation cascade reactions depend on the calcium present in the plasma, not the platelet activation step itself. Therefore, platelets may very well become activated in citrate-anticoagulated blood (e.g., as a result of poor handling, blood drawing techniques, or prolonged storage). Nevertheless, it needs also to be remembered that the activation processes in the absence of extracellular calcium are not complete—especially if an ionophore is used as a stimulant [106].

The topic of platelet activation by anticoagulants deserves a separate mention. In particular, heparin was shown to activate platelets at least under some conditions [31, 40].

Hydrolysis of PIP<sub>2</sub> also generates DAG (green hexagons in Fig. 3), which activates some of the protein kinase C (PKC) isoforms. Other isoforms are activated by Ca, yet others require both of these co-factors, while some of the more recently discovered isoforms depend on other factors for activation [49].

An important feature of the platelet signalling network appears to be the incorporation of extracellular events (such as secretion and aggregation) into the signalling pathways. This can be illustrated by following the effect of ADP (blue solid arrows in Fig. 3). Its binding to the P<sub>2</sub>Y<sub>1</sub> receptor leads to the activation of PLC $\beta$  through the G<sub>q</sub> protein and a transient rise in the intracellular calcium level through the SOCE mechanism, which drives platelet shape change. Its binding to the P<sub>2</sub>Y<sub>12</sub> receptor drives GPIIb/IIIa conformational change. Platelet aggregation, which is initially reversible, [90] leads to the activation of PLA<sub>2</sub> and thromboxane synthesis. TXA<sub>2</sub> is released by the platelets (purple dashed arrow in Fig. 3) and interacts with own receptor, leading to the dense granule secretion and other responses (solid purple arrows in Fig. 3). Aggregation becomes irreversible [90]. Aspirin, which blocks TXA<sub>2</sub> production, inhibits dense granule secretion and the second (irreversible) phase of aggregation, but not the calcium rise, the shape change, or the first (reversible) wave of aggregation caused by ADP (Fig. 2 in ref. [103]). In the absence of aggregation, ADP does not lead to thromboxane production or dense granule secretion [12]. On the other hand, aspirin had no effect on the expression of P-selectin or the secretion of  $\beta$ -thromboglobulin (both  $\alpha$ -granule markers) by platelets stimulated with ADP in plasma, while inhibiting irreversible aggregation and dense granule secretion as measured by ADP and serotonin release [99, 101]. We also observed P-selectin expression in purified platelets stimulated with ADP [45].

Staged activation of platelets can also be illustrated with other examples. ADP is involved in activating the GPIa/IIa (integrin  $\alpha_{IIb}\beta_1$ ) [64]. This process allows it to bind collagen with high affinity at low exposure levels (of collagen). Collagen then stimulates GPVI receptor, leading to other responses such as TXA<sub>2</sub> production and PS expression (dashed brown arrow in Fig. 3) [51].

Similarly, ADP is involved in the transition from reversible to irreversible aggregates in platelets activated by vWF at high shear [83]. The signalling pathway from the vWF receptor to the secretion of dense granules is illustrated in Fig. 3 with a solid brown arrow. (The effect of aspirin was not investigated in that study, but unless GPIb/vWF interaction simultaneously leads to ADP secretion and TXA<sub>2</sub> production, a three-stage process is actually involved: GPIb/vWF leading to ADP secretion which in turn leads to TXA<sub>2</sub> production as described above). Furthermore, a similar effect is observed in the case of a non-physiological agonist, phorbol 12-myristate 13-acetate (PMA). PMA does not act via a receptor but directly activates protein kinase C (PKC), located downstream of the PLC $\beta$ -dependent calcium rise. It appears that the elevation of the intracellular calcium in PMA-treated platelets occur via an autocrine mechanism involving the secretion dense granules [48].



Such “staging” of the activation process appears to be a general feature of the platelet signalling pathways, as is the integration of extracellular events into the signalling pathways. This strategy serves an amplification function but also provides a key control mechanism for limiting platelet activation. One practical consequence of this is that platelet signalling is best studied with purified platelets, because various components of plasma may complete the signalling pathways without the control or knowledge of the investigator.

Ultimately, platelet signalling network funnels all of the incoming signals towards responses such as shape change, GPIIb/IIIa activation (and consequent aggregation), PS expression, and granule secretion through the amplifying autocrine steps and synergistic pathways. The response to each individual agonist, however, is different. Some of the differences between ADP and other agonists have already been illustrated above, as were some of the TXA<sub>2</sub>-specific effects. Differences in responses to individual agonists arise through several mechanisms. On one hand, there is evidence of different threshold calcium concentrations required for the different responses, such as shape changes, aggregation, and secretion [106]. In that study, intracellular calcium levels were measured with Ca-sensitive fluorescent dyes in platelets treated with various agonists and a calcium ionophore ionomycin. The same study showed lower intracellular calcium levels accompanying responses in platelets treated with thrombin, implying the existence of Ca-independent signalling pathways. For detailed discussion of calcium-independent platelet activation pathways, the reader is referred to refs. [98, 102–105]. The two sets of pathways—calcium dependent and calcium-independent—act in synergy with each other.

On the other hand, the temporal profile of the calcium rise also plays a role. A well-documented example of this is the difference between PAR1 and PAR4 thrombin receptors in terms of the timing and duration of the calcium rise and in terms of the functions the two receptors play in early and late stages of thrombus formation [3, 18, 22, 128].

Ultimately, a platelet is akin to a piano (Fig. 4): its response is both carefully timed and spatially orchestrated to achieve a specific “melody” of the plug formation, clot formation, clot retraction, and subsequent wound healing processes, each initiated at the right location of the wound site and at the right time. This choreographed sequence of events is revealed, for example, at the level of the blood clot organization, with differences in platelet activation, packing density, fibrin formation, and transport of material through the clot, in the different areas of the clot [86, 126]. Some exciting work is now being done in this area, also in terms of linking clot organization to the way platelet signalling network integrates information from the agonist receptors [86, 126]. Indeed, differences in responses to individual agonists and their synergies hold the key to the spatiotemporal organization of the platelet response [86]. They also offer potential points for selective intervention, exploited by such agents as aspirin and P2Y<sub>12</sub> receptor antagonists [60].

This kind of organization presupposes the ability of the signalling network elements to integrate the incoming signals over a period of time and over the different input channels (receptors for different agonists). Much remains to be learned about this sort of signal integration in the regulation of platelet activity even in the context of their haemostatic function, but especially—concerning the integration of cytokine receptors, pattern-matching toll-like receptors for microbial recognition, interactions with other cells, and other processes concerned with non-haemostatic platelet functions, [15, 107, 121, 148] into the signalling network.

To illustrate some of the limitations of the current approaches for studying differences between signalling pathways of the different agonists, we describe what is known about the secretion of  $\alpha$ - and dense granules. These two sets of granules are different morphologically and in terms of their contents. Logically, the machinery that regulates their secretion also appears to be different: there are reports of




Fig. 4. Spectrum of platelet responses to various stimuli. This figure is meant to illustrate that platelet activation by different agonists is subtly different. The platelet signalling pathways are likened to a piano, with each agonist eliciting a unique “chord” of a response. Some of the keys overlap between agonists, but some are different. In this way, various combinations of agonists produce the melody of platelet responses. Platelets are depicted as grey circles, as in Fig. 1. Differences between platelets are illustrated in terms of different expression levels of CD62P (red circles) and CD63 (yellow circles) on platelets stimulated by thrombin and ADP, and in terms of differences in platelet surface glycosylation. The latter are based on our recent work on platelet surface glycosylation and its changes caused by agonist treatment [43, 45]. For example, thrombin activated platelets are shown to exhibit higher levels of expression of N-acetyl glucoseamine (blue squares) and lower levels of sialic acid (purple rhombi) than ADP-stimulated ones. N-acetyl glucoseamine is also shown to cluster on the thrombin-activated platelets. This figure is intended only as an illustration of the finding that platelets express different sets of carbohydrates, depending on how they were stimulated. The inspiration for this figure came from the piano analogy used by J. Tepperman and H. M. Tepperman in their endocrinology text to describe the role of sympathetic nervous system and adrenal medulla in the fight-or-flight response (Tepperman J, Tepperman HM. *Metabolic and endocrine physiology: an introductory text*. Year Book Medical Publishers, Chicago, 1987 (fifth edition).

the GTPase Rab4 being involved in regulating the  $\alpha$ -granule secretion, and of Rab27b—in the secretion of dense granules [122, 131, 149]. There are several examples of differential secretion of  $\alpha$ - and dense granules: [9, 28, 44, 66, 130].

- Investigating the effect of cytoskeleton rearrangements on granule secretion Flaumenhaft et al. [28] observed opposite effects: high concentrations (above  $\sim 100$   $\mu$ M) of latrunculin A, which facilitates actin cytoskeleton disassembly, inhibited  $\alpha$ -granule secretion but stimulate dense granule secretion in TRAP- or PMA-stimulated platelets. That study was done on platelets purified by gel filtration from citrate anticoagulated blood. Flow cytometry was used to monitor the expression of  $\alpha$ -granule marker CD62P. Dense granule secretion was monitored by measuring levels of ATP and serotonin release.
- Taylor et al. [130] using citrate-anticoagulated platelet-rich plasma and flow cytometry assays for CD62P and CD63 expression examined the dependence of granule release on the type and concentration of various agonists. They found that the kinetics of the expression of the two markers was different and granule secretion events that did not correlate with platelet aggregation responses.
- Broberg et al. [9] using citrate anticoagulated platelet-rich plasma, identified surface-adsorbed vWF as a specific regulator of CD62P expression on adhering platelets; adsorbed vWF did not have such an effect on the release of other markers used in that study (PF4 and ATP for  $\alpha$ -granules and dense granules, respectively). This study is noteworthy because it also indicates that CD62P expression and secretion of  $\alpha$ -granule contents are not always correlated.
- Kang et al., [66] studying adhesion of platelets to surfaces of various peptides from platelet-rich plasma and whole blood, showed that serotonin release (dense granules) correlated with the number of adhering platelets but  $\beta$ -thromboglobulin release ( $\alpha$ -granules) exhibited a maximum for a particular set of peptides. Unfortunately, in their study, the  $\alpha$ - and dense granule release were not compared under identical conditions.
- In our own study of platelets, purified from citrate-anticoagulated blood by centrifugation, adhering on TiO<sub>2</sub> in the absence of extracellular calcium, we found that CD63 but not CD62P was expressed [44].

Although it is clear that secretion of these two sets of granules is regulated by separate pathways, [144] neither the physiological significance, nor the physiologically relevant triggers of such differential behaviour are known; individual signalling pathways connecting agonists to the secretion of each set of granules have also not been identified—and this is despite the fact that markers for the secretion of each of the sets of granules have been readily available for quite some time.

#### **4. Need for new markers and functional assays to monitor platelet activation**

The limitations of the existing repertoire of markers and assays for studying platelet activation are apparent already at the level of platelet haemostatic function. Considering their other functions, the situation grows more complex and the need for new assays and markers for studying platelets becomes more acute.

A reasonable place to start developing such assays is  $\alpha$ -granule secretion. Given that  $\alpha$ -granules contain active substances with contradictory functions, it is logical to suppose that their secretion is thematically controlled. Initial reports by Italiano et al. appeared to support this idea. These authors reported that pro-angiogenic vascular endothelial growth factor (VEGF) and thrombospondin-1 were packaged into a

different population of  $\alpha$ -granules than the anti-angiogenic endostatin and basic fibroblast growth factor. Fibrinogen and von Willebrand factor (vWF) were also found in distinct  $\alpha$ -granule subpopulations. Moreover, these authors also reported their thematic secretion in response to stimulation with different agonists [59].

Cryo-electron tomography confirms the notion, that different kinds of  $\alpha$ -granules exist within the platelets [134]. It also shows that vWF and fibrinogen are localized to different granule subtypes. Another, earlier observation of differences in granule motion in adhering platelets that correlated with the distribution of vesicle-associated membrane protein (VAMP) isoforms, similarly confirms the notion that there are different  $\alpha$ -granule subpopulations in the platelets [94].

However, other conclusions of Italiano have been challenged. Peters et al. [94] noted that VEGF co-localized with different VAMP isoforms (thought to be responsible for the secretion of different granule subpopulations). In a study of 15 different  $\alpha$ -granule proteins, Kamykowski et al. reported no clustering patterns of pro- vs. anti-angiogenic proteins [65]. They did report that fibrinogen and vWF were spatially segregated, but concluded that  $\alpha$ -granule content could be organized in different partitions, or zones, within the granules. Subsequently, Rönnuland et al. used super-resolution fluorescence microscopy (STED; one of the techniques the developers of which received a Nobel Prize in Chemistry in 2014) to examine the distribution of various  $\alpha$ -granule cargo. They showed no evidence of thematic packaging, [108] suggesting that the resolution of confocal microscopy was not sufficient to discern such patterns. In the same study, Rönnuland et al. also showed that granule contents was organized into structures with sizes much smaller than the granules themselves, suggesting sub-granule zones, and that treating platelets with different agonists lead to distinctly different distributions of fluorescently labelled molecules. However, Jonnalagadda et al., analysing the release of 28 molecules known to reside in the  $\alpha$ -granules by micro-ELISA arrays from platelet stimulated with different agonists as a function of time, did not observe agonist-specific release patterns. They concluded that the release was stochastic rather than thematic, and that agonist strength and kinetic effects played an important role in determining platelet secretion profile [63].

This survey of the latest attempts to identify markers of distinct platelet functional states through differential  $\alpha$ -granule secretion shows, that this approach has so far not met with success. With the goal of expanding the range of markers that could be used to study various aspects of platelet physiology, we have recently examined changes in platelet surface glycosylation induced by treating platelets with various physiological and non-physiological agonists [43, 45]. The surfaces of platelets, like surfaces of all cells, are covered with carbohydrates. Most of them originate from cell surface glycoproteins [87]. Surprisingly little is known about the platelet carbohydrate coat, and even less is known about how it changes upon platelet activation:

- Structures of the carbohydrates linked to several of the surface glycoproteins have been elucidated [71, 72, 132, 133].
- Carbohydrate moieties on the GPVI complex are thought to be involved in its interactions with collagen [74].
- The chemical nature of the platelet carbohydrate coat changes as platelets age [140]. These changes serve as signals for platelet clearance by the liver macrophages and hepatocytes, [53, 111, 124] by mechanisms similar to those involved in clearance of aged erythrocytes and glycosylated plasma proteins [2].
- Carbohydrate-binding proteins selectins [1, 20] are present on platelet, leukocyte, and endothelial cell surfaces. They mediate adhesion and rolling of leukocytes, and CD62P – P-selectin—is involved in the platelet-leukocyte and platelet-endothelial cell interactions. This topic has already been

addressed in the very beginning of this article, when platelet role in maintaining vascular integrity was discussed [67, 139]. Selectins belong to the same family of the carbohydrate binding proteins (C-type lectins [1]) as the hepatic asialoglycoprotein receptor (ASGPR), already mentioned in the context of platelet clearance [41, 112].

- Galectins, [19] members of another lectin family ( $\beta$ -Galactose-binding proteins), interact with platelet surface carbohydrates and act as platelet agonists *in vitro*, but their *in vivo* role is not clear [113].
- Platelets release glycosyltransferases and regulate their activity as a part of the activation process, [140] but the role or the mechanisms of this regulation in the context of platelet activation are not understood.
- Differences in the composition of the platelet carbohydrate coat between platelets from healthy individuals and those from uremic or diabetic patients have been reported [81, 82].
- While progress is being made systematically analysing the glycome of, e.g., the immune system [50] and vascular endothelium, [120] a literature search using keywords “glycome” and “platelets” returns six publications, two of which are meeting abstracts. Similarly, among the entries found on the website of the Consortium for Functional Genomics (<http://www.functionalglycomics.org/static/index.shtml>), only one involves platelets [56]. In this context, a very recent review of the role surface carbohydrates play in platelet-endothelial cell interactions should also be mentioned [26].

Recently, we used fluorescently labelled plant lectins [1] to investigate platelet surface glycosylation changes as a result of agonist treatment. Using a panel of nine lectins and seven different agonists, we detected changes in platelet surface glycosylation upon stimulation and showed that these changes were agonist-specific (see also the illustration in Fig. 4) [45]. Some of them could be correlated with the expression of the granule-specific glycoproteins (CD62P for the  $\alpha$ -granules and CD63 for the dense granules), while other changes did not correlate with secretion events. Our results allowed us to conclude that differences in the individual signalling pathways between different agonists were responsible for the differences in lectin binding to platelets activated with these agonists. While many questions remain to be answered, our study does suggest monitoring platelet surface glycosylation changes as a promising approach to studying platelet activation in the laboratory and to developing clinical diagnostic assays. It certainly offers much more in terms of diversity than existing antibody or soluble markers.

## 5. Development of assays for studying non-haemostatic platelet functions

While the development of new labels for studying platelet activation is a necessary step for unravelling their physiology and providing quantitative data for systems-biology approaches to model their signalling networks, all of the above studies share the same fundamental flaw: they focus on the range of agonists that stimulate procoagulant platelet functions. In order to understand the diversity of platelet functions, *in vitro* model systems need to be developed that transcend the reliance on the traditional set of platelet agonists. Possible search directions include co-cultures with endothelial cells and mesenchymal stem cells; interactions with leukocytes; use of inflammatory mediators as agonists or functional modulators; interactions with implants—because in this case, platelets are subjected to activating conditions that differ significantly from the natural ones they encounter in the body, and this may offer a unique insight into their physiology. Such a conceptual switch from the “coagulation” frame of mind is likely to lead to breakthroughs in the understanding of platelet regulation that will translate into their use in personalized, regenerative therapies. Indeed, such approaches are being explored [25].

## Acknowledgments

The author wishes to thank Dr. Swati Gupta (CIC biomaGUNE, San Sebastian, Spain) for her help with designing Fig. 3 and numerous discussions we had on the subject of platelet activation and signalling. Financial support of the Biointerfaces program (BIF) of the Helmholtz Association at the Karlsruhe Institute of Technology (KIT) is also gratefully acknowledged.

## References

- [1] M. Ambrosi, N.R. Cameron and B.G. Davis, Lectins: Tools for the molecular understanding of the glycodecode, *Org Biomol Chem* **3** (2005), 1593–1608.
- [2] D. Aminoff, W.F. Bruegge, W.C. Bell, K. Sarpolis and R. Williams, Role of sialic acid in survival of erythrocytes in the circulation: Interaction of neuraminidase-treated and untreated erythrocytes with spleen and liver at the cellular level, *Proc Natl Acad Sci U S A* **74** (1977), 1521–1524.
- [3] H. Andersen, D.L. Greenberg, K. Fujikawa, W. Xu, D.W. Chung and E.W. Davie, Protease-activated receptor 1 is the primary mediator of thrombin-stimulated platelet procoagulant activity, *Proc Natl Acad Sci U S A* **96** (1999), 11189–11193.
- [4] E. Anitua, Plasma rich in growth factors: Preliminary results of use in the preparation of future sites for implants, *Int J Oral Maxillofac Implants* **14** (1999), 529–535.
- [5] E. Anitua, I. Andia, B. Ardanza, P. Nurden and A.T. Nurden, Autologous platelets as a source of proteins for healing and tissue regeneration, *Thromb Haemost* **91** (2004), 4–15. 10.1160/th03-07-0440.
- [6] E. Anitua, M. Sanchez, R. Prado and G. Orive, Plasma rich in growth factors: The pioneering autologous technology for tissue regeneration, *Journal of biomedical materials research* **97** (2011), 536.
- [7] K. Balasubramanian and A.J. Schroit, Aminophospholipid asymmetry: A matter of life and death, *Annu Rev Physiol* **65** (2003), 701–734.
- [8] D.L. Bhatt and E.J. Topol, Scientific and therapeutic advances in antiplatelet therapy, *Nature Reviews Drug Discovery* **2** (2003), 15–28. 10.1038/nrd985.
- [9] M. Broberg and H. Nygren, Von Willebrand factor, a key protein in the exposure of CD62P on platelets, *Biomaterials* **22** (2001), 2403–2409.
- [10] T. Burnouf, H.A. Goubran, T.M. Chen, K.L. Ou, M. El-Ekiaby and M. Radosevic, Blood-derived biomaterials and platelet growth factors in regenerative medicine, *Blood Rev* **27** (2013), 77–89. 10.1016/j.blre.2013.02.001.
- [11] M. Cattaneo, Resistance to anti-platelet agents, *Thrombosis Research* **127** (Suppl 3) (2011), S61–S63. 10.1016/S0049-3848(11)70017-2.
- [12] M. Cattaneo, C. Gachet, J.P. Cazenave and M.A. Packham, Adenosine diphosphate (ADP) does not induce thromboxane A<sub>2</sub> generation in human platelets, *Blood* **99** (2002), 3868–3869; author reply 9–70.
- [13] J. Choi and J.C. Kermode, New therapeutic approaches to combat arterial thrombosis: Better drugs for old targets, novel targets, and future prospects, *Molecular Interventions* **11** (2011), 111–123. 10.1124/mi.11.2.9.
- [14] G. Cimmino and P. Golino, Platelet biology and receptor pathways, *Journal of Cardiovascular Translational Research* **6** (2013), 299–309. 10.1007/s12265-012-9445-9.
- [15] F. Cognasse, H. Hamzeh-Cognasse, S. Lafarge, O. Delezay, B. Pozzetto, A. McNicol and O. Garraud, Toll-like receptor 4 ligand can differentially modulate the release of cytokines by human platelets, *British Journal of Haematology* **141** (2008), 84–91. 10.1111/j.1365-2141.2008.06999.x.
- [16] B.S. Coller, Historical perspective and future directions in platelet research, *Journal of Thrombosis and Haemostasis: JTH* **9** (Suppl 1) (2011), 374–395. 10.1111/j.1538-7836.2011.04356.x.
- [17] S.R. Coughlin, Thrombin signalling and protease-activated receptors, *Nature* **407** (2000), 258–264. 10.1038/35025229.
- [18] L. Covic, A.L. Gresser and A. Kuliopulos, Biphasic kinetics of activation and signaling for PAR1 and PAR4 thrombin receptors in platelets, *Biochemistry* **39** (2000), 5458–5467.
- [19] R.D. Cummings and F.T. Liu, Galectins. In: A. Varki, R.D. Cummings, J.D. Esko, H.H. Freeze, P. Stanley, C.R. Bertozzi, G.W. Hart, M.E. Etzler, eds. *Essentials of Glycobiology*, 2nd edn. Cold Spring Harbor, NY, 2009.
- [20] R.D. Cummings and R.P. McEver, C-type Lectins. In: A. Varki, R.D. Cummings, J.D. Esko, H.H. Freeze, P. Stanley, C.R. Bertozzi, G.W. Hart, M.E. Etzler, eds. *Essentials of Glycobiology*, 2nd edn. Cold Spring Harbor, NY, 2009.
- [21] J.F. Danielli, Capillary permeability and oedema in the perfused frog, *The Journal of physiology* **98** (1940), 109–129.

- [22] E. De Candia, Mechanisms of platelet activation by thrombin: A short history, *Thrombosis Research* **129** (2012), 250–256. 10.1016/j.thromres.2011.11.001.
- [23] C.A. Demopoulos, R.N. Pinckard and D.J. Hanahan, Platelet-activating factor - evidence for 1-o-alkyl-2-acetyl-sn-glycerol-3-phosphorylcholine as the active component (a new class of lipid chemical mediators), *J Biol Chem* **254** (1979), 9355–9358.
- [24] B.L. Eppley, W.S. Pietrzak and M. Blanton, Platelet-rich plasma: A review of biology and applications in plastic surgery, *Plast Reconstr Surg* **118** (2006), 147E–159E. 10.1097/01.prs.0000239606.92676.cf.
- [25] J. Etulain, C. Fondevila, S. Negrotto and M. Schattner, Platelet-mediated angiogenesis is independent of VEGF and fully inhibited by aspirin, *British Journal of Pharmacology* **170** (2013), 255–265. 10.1111/bph.12250.
- [26] J. Etulain and M. Schattner, Glycobiology of platelet-endothelial cell interactions, *Glycobiology* **24** (2014), 1252–1259. 10.1093/glycob/cwu056.
- [27] R. Fitzgerald and M. Pirmohamed, Aspirin resistance: Effect of clinical, biochemical and genetic factors, *Pharmacology & Therapeutics* **130** (2011), 213–225. 10.1016/j.pharmthera.2011.01.011.
- [28] R. Flaumenhaft, J.R. Dilks, N. Rozenvayn, R.A. Monahan-Earley, D. Feng and A.M. Dvorak, The actin cytoskeleton differentially regulates platelet alpha-granule and dense-granule secretion, *Blood* **105** (2005), 3879–3887. 10.1182/blood-2004-04-1392.
- [29] P.S. Frenette, R.C. Johnson, R.O. Hynes and D.D. Wagner, Platelets roll on stimulated endothelium *in vivo*: An interaction mediated by endothelial P-selectin, *Proc Natl Acad Sci U S A* **92** (1995), 7450–7454.
- [30] A. Gaarder, J. Jonsen, S. Laland, A. Hellem and P.A. Owren, Adenosine diphosphate in red cells as a factor in the adhesiveness of human blood platelets, *Nature* **192** (1961), 531–532.
- [31] C. Gao, B. Boylan, J. Fang, D.A. Wilcox, D.K. Newman and P.J. Newman, Heparin promotes platelet responsiveness by potentiating alphaIIb beta3-mediated outside-in signaling, *Blood* **117** (2011), 4946–4952.
- [32] A.E. Garcia, S.R. Mada, M.C. Rico, R.A. Dela Cadena and S.P. Kunapuli, Clopidogrel, a P2Y12 receptor antagonist, potentiates the inflammatory response in a rat model of peptidoglycan polysaccharide-induced arthritis, *PLoS One* **6** (2011), e26035. 10.1371/journal.pone.0026035.
- [33] G.J. Gasic, T.B. Gasic and C.C. Stewart, Antimetastatic effects associated with platelet reduction, *Proc Natl Acad Sci U S A* **61** (1968), 46–52.
- [34] L.J. Gay and B. Felding-Habermann, Contribution of platelets to tumour metastasis, *Nature Reviews Cancer* **11** (2012), 123–134. 10.1038/nrc3004.
- [35] P.L.A. Giesen, G.M. Willems and W.T. Hermens, Production of thrombin by the prothrombinase complex is regulated by membrane-mediated transport of prothrombin, *J Biol Chem* **266** (1991), 1379–1382.
- [36] M.A. Gimbrone Jr., R.H. Aster, R.S. Cotran, J. Corkery, J.H. Jandl and J. Folkman, Preservation of vascular integrity in organs perfused *in vitro* with a platelet-rich medium, *Nature* **222** (1969), 33–36.
- [37] T. Goerge, B. Ho-Tin-Noe, C. Carbo, C. Benarafa, E. Remold-O'Donnell, B.Q. Zhao, S.M. Cifuni and D.D. Wagner, Inflammation induces hemorrhage in thrombocytopenia, *Blood* **111** (2008), 4958–4964. 10.1182/blood-2007-11-123620.
- [38] J. Golanski, T. Pietrucha, Z. Baj, J. Greger and C. Watala, Molecular insights into the anticoagulant-induced spontaneous activation of platelets in whole blood-various anticoagulants are not equal, *Thromb Res* **83** (1996), 199–216.
- [39] M.B. Gorbet and M.V. Sefton, Biomaterial-associated thrombosis: Roles of coagulation factors, complement, platelets and leukocytes, *Biomaterials* **25** (2004), 5681–5703.
- [40] A. Greinacher, Platelet activation by heparin, *Blood* **117** (2011), 4686–4687.
- [41] P.K. Grewal, The Ashwell-Morell receptor, *Methods in Enzymology* **479** (2010), 223–241. 10.1016/S0076-6879(10)79013-3.
- [42] Y.L. Gu, M.L. Fokkema, M.A. Kampinga, B.J. de Smet, E.S. Tan, A.F. van den Heuvel and F. Zijlstra, Intracoronary versus intravenous abciximab in ST-segment elevation myocardial infarction: Rationale and design of the CICERO trial in patients undergoing primary percutaneous coronary intervention with thrombus aspiration, *Trials* **10** (2009), 90. 10.1186/1745-6215-10-90.
- [43] S. Gupta, Selective Activation of Platelets by Surfaces and Soluble Agonists. *Department of Biochemistry and Molecular Biology*, University of the Basque Country, Leioa, 2014.
- [44] S. Gupta and I. Reviakine, Platelet activation profiles on TiO<sub>2</sub>: Effect of Ca<sup>2+</sup> binding to the surface, *Biointerphases* **7** (2012), 28–40.
- [45] S. Gupta and I. Reviakine, The sweeter aspects of platelet activation: A lectin-based assay reveals agonist-specific glycosylation patterns, *Biochim Biophys Acta* **1840** (2014), 3423–3433.



- [46] A.V. Gyulkhandanyan, A. Mutlu, J. Freedman and V. Leytin, Selective triggering of platelet apoptosis, platelet activation or both, *British Journal of Haematology* **161** (2013), 245–254. 10.1111/bjh.12237.
- [47] I.A. Hagberg and T. Lyberg, Blood platelet activation evaluated by flow cytometry: Optimised methods for clinical studies, *Platelets* **11** (2000), 137–150.
- [48] M.T. Harper, M.J. Mason, S.O. Sage and A.G. Harper, Phorbol ester-evoked Ca<sup>2+</sup> signaling in human platelets is via autocrine activation of P(2X1) receptors, not a novel non-capacitative Ca<sup>2+</sup> entry, *Journal of Thrombosis and Haemostasis: JTH* **8** (2010), 1604–1613. 10.1111/j.1538-7836.2010.03867.x.
- [49] M.T. Harper and A.W. Poole, Diverse functions of protein kinase C isoforms in platelet activation and thrombus formation, *J Thromb Haemost* **8** (2010), 454–462. JTH3722 [pii] 10.1111/j.1538-7836.2009.03722.x.
- [50] S.M. Haslam, S. Julien, J.M. Burchell, C.R. Monk, A. Ceroni, O.A. Garden and A. Dell, Characterizing the glycome of the mammalian immune system, *Immunol Cell Biol* **86** (2008), 564–573.
- [51] J.W. Heemskerk, W.M. Vuist, M.A. Feijge, C.P. Reutelingsperger and T. Lindhout, Collagen but not fibrinogen surfaces induce bleb formation, exposure of phosphatidylserine, and procoagulant activity of adherent platelets: Evidence for regulation by protein tyrosine kinase-dependent Ca<sup>2+</sup> responses, *Blood* **90** (1997), 2615–2625.
- [52] P.M. Henson and R.N. Pinckard, Platelet activating factor (PAF), A possible direct mediator of anaphylaxis in the rabbit and a trigger for the vascular deposition of circulating immune complexes, *Monographs in Allergy* **12** (1977), 13–26.
- [53] K.M. Hoffmeister, E.C. Josefsson, N.A. Isaac, H. Clausen, J.H. Hartwig and T.P. Stossel, Glycosylation restores survival of chilled blood platelets, *Science (New York, NY)* **301** (2003), 1531–1534.
- [54] D.R. Holmes Jr., D.J. Kereiakes, N.S. Kleiman, D.J. Moliterno, G. Patti and C.L. Grines, Combining antiplatelet and anticoagulant therapies, *Journal of the American College of Cardiology* **54** (2009), 95–109. 10.1016/j.jacc.2009.03.044.
- [55] B. Ho-Tin-Noe, M. Demers and D.D. Wagner, How platelets safeguard vascular integrity, *Journal of Thrombosis and Haemostasis: JTH* **9**(Suppl 1) (2011), 56–65. 10.1111/j.1538-7836.2011.04317.x.
- [56] H. Hu, P.C. Armstrong, E. Khalil, Y.C. Chen, A. Straub, M. Li, J. Soosairajah, C.E. Hagemeyer, N. Bassler, D. Huang, I. Ahrens, G. Krippner, E. Gardiner and K. Peter, GPVI and GPIIb/IIIa mediate staphylococcal superantigen-like protein 5 (SSL5) induced platelet activation and direct toward glycans as potential inhibitors, *PLoS One* **6** (2011), e19190. 10.1371/journal.pone.0019190.
- [57] H. Hu, M. Forslund and N. Li, Influence of extracellular calcium on single platelet activation as measured by whole blood flow cytometry, *Thromb Res* **116** (2005), 241–247.
- [58] Y. Ikeda, M. Murata and S. Goto, *von Willebrand factor-dependent shear-induced platelet aggregation: Basic mechanisms and clinical implications*. In: F. Numano and R. Ross, eds. *Atherosclerosis IV: Recent Advances in Atherosclerosis Research: The Fourth Saratoga International Conference on Atherosclerosis*. New York: New York Acad Sciences, 1997, pp. 325–336.
- [59] J.E. Italiano Jr., J.L. Richardson, S. Patel-Hett, E. Battinelli, A. Zaslavsky, S. Short, S. Ryeom, J. Folkman and G.L. Klement, Angiogenesis is regulated by a novel mechanism: Pro- and antiangiogenic proteins are organized into separate platelet alpha granules and differentially released, *Blood* **111** (2008), 1227–1233.
- [60] S.P. Jackson and S.M. Schoenwaelder, Antiplatelet therapy: In search of the ‘magic bullet’, *Nat Rev Drug Discov* **2** (2003), 775–789.
- [61] P.A. Janmey, J.P. Winer and J.W. Weisel, Fibrin gels and their clinical and bioengineering applications, *Journal of the Royal Society Interface* **6** (2009), 1–10. 10.1098/rsif.2008.0327.
- [62] J. Jin and S.P. Kunapuli, Coactivation of two different G protein-coupled receptors is essential for ADP-induced platelet aggregation, *Proc Natl Acad Sci U S A* **95** (1998), 8070–8074.
- [63] D. Jonnalagadda, L.T. Izu and S.W. Whiteheart, Platelet secretion is kinetically heterogeneous in an agonist-responsive manner, *Blood* **120** (2011), 5209–5216.
- [64] S.M. Jung and M. Moroi, Platelet collagen receptor integrin alpha2beta1 activation involves differential participation of ADP-receptor subtypes P2Y1 and P2Y12 but not intracellular calcium change, *European Journal of Biochemistry/FEBS* **268** (2001), 3513–3522.
- [65] J. Kamykowski, P. Carlton, S. Sehgal and B. Storrie, Quantitative immunofluorescence mapping reveals little functional coclustering of proteins within platelet alpha-granules, *Blood* **118** (2011), 1370–1373.
- [66] I.K. Kang, Y. Ito, M. Sisido and Y. Imanishi, Serotonin and beta-thromboglobulin release reaction from platelet as triggered by interaction with polypeptide derivatives, *J Biomed Mater Res* **22** (1988), 595–611.
- [67] Z.S. Kaplan and S.P. Jackson, The role of platelets in atherothrombosis, *Hematology Am Soc Hematol Educ Program* **2011** (2011), 51–61.

- [68] L.R. Kark, J.M. Karp and J.E. Davies, Platelet releasate increases the proliferation and migration of bone marrow-derived cells cultured under osteogenic conditions, *Clinical Oral Implants Research* **17** (2006), 321–327. 10.1111/j.1600-0501.2005.01189.x.
- [69] B. Kehrel, S. Wierwille, K.J. Clemetson, O. Anders, M. Steiner, C.G. Knight, R.W. Farndale, M. Okuma and M.J. Barnes, Glycoprotein VI is a major collagen receptor for platelet activation: It recognizes the platelet-activating quaternary structure of collagen, whereas CD36, glycoprotein IIb/IIIa, and von Willebrand factor do not, *Blood* **91** (1998), 491–499.
- [70] A.A. Khorana, Malignancy, thrombosis and Trousseau: The case for an eponym, *Journal of Thrombosis and Haemostasis: JTH* **1** (2003), 2463–2465.
- [71] S.A. Korrel, K.J. Clemetson, H. van Halbeek, J.P. Kamerling, J.J. Sixma and J.F. Vliegthart, Identification of a tetrasialylated monofucosylated tetraantennary N-linked carbohydrate chain in human platelet glycolalicin, *FEBS Letters* **228** (1988), 321–326.
- [72] S.A. Korrel, K.J. Clemetson, H. Van Halbeek, J.P. Kamerling, J.J. Sixma and J.F. Vliegthart, Structural studies on the O-linked carbohydrate chains of human platelet glycolalicin, *European Journal of Biochemistry/FEBS* **140** (1984), 571–576.
- [73] A. Kruger, C. Mrowietz, A. Lendlein and F. Jung, Interaction of human umbilical vein endothelial cells (HUVEC) with platelets *in vitro*: Influence of platelet concentration and reactivity, *Clinical Hemorheology and Microcirculation* **55** (2013), 111–120. 10.3233/CH-131695.
- [74] T.J. Kunicki, Y. Cheli, M. Moroi and K. Furihata, The influence of N-linked glycosylation on the function of platelet glycoprotein VI, *Blood* **106** (2005), 2744–2749. 10.1182/blood-2005-04-1454.
- [75] M. Leslie, Beyond clotting: The powers of platelets, *Science* (New York, NY) **328** (2010), 562–564.
- [76] V. Leytin, Apoptosis in the anucleate platelet, *Blood Rev* **26** (2012), 51–63. 10.1016/j.blre.2011.10.002.
- [77] Z. Li, M.K. Delaney, K.A. O'Brien and X. Du, Signaling during platelet adhesion and activation, *Arterioscler Thromb Vasc Biol* **30** (2011), 2341–2349.
- [78] P. Libby, P.M. Ridker and G.K. Hansson, Progress and challenges in translating the biology of atherosclerosis, *Nature* **473** (2011), 317–325. 10.1038/nature10146.
- [79] A.M. Lincoff, D.J. Kereiakes, M.A. Mascelli, L.I. Deckelbaum, E.S. Barnathan, K.K. Patel, B. Frederick, M.T. Nakada and E.J. Topol, Abciximab suppresses the rise in levels of circulating inflammatory markers after percutaneous coronary revascularization, *Circulation* **104** (2001), 163–167.
- [80] F. Lombardi, C. De Chaumont, D.C. Shields and N. Moran, Platelet signalling networks: Pathway perturbation demonstrates differential sensitivity of ADP secretion and fibrinogen binding, *Platelets* **23** (2012), 17–25. 10.3109/09537104.2011.594190.
- [81] C. Lupu, M. Rizescu and M. Calb, Altered distribution of some surface glycosaminoglycans and glycoconjugates on human blood-platelets in diabetes-mellitus, *Platelets* **5** (1994), 201–208. 10.3109/09537109409006047.
- [82] M. Manso, I. De Dios, L. Alberca and V. Vicente, Studies on platelet surface carbohydrates in normal and uraemic platelets using 125I-labelled lectins, *Blut* **50** (1985), 287–292.
- [83] M.J. Maxwell, E. Westein, W.S. Nesbitt, S. Giuliano, S.M. Dopheide and S.P. Jackson, Identification of a 2-stage platelet aggregation process mediating shear-dependent thrombus formation, *Blood* **109** (2007), 566–576. 10.1182/blood-2006-07-028282.
- [84] A.D. Michelson, *Platelets*, Academic Press, London, 2012.
- [85] D.M. Monroe, M. Hoffman and H.R. Roberts, Platelets and thrombin generation, *Arteriosclerosis, Thrombosis, and Vascular Biology* **22** (2002), 1381–1389.
- [86] I.C. Munnix, J.M. Cosemans, J.M. Auger and J.W. Heemskerk, Platelet response heterogeneity in thrombus formation, *Thromb Haemost* **102** (2009), 1149–1156. 10.1160/TH09-05-0289.
- [87] A.T. Nurden, Platelet membrane glycoproteins: A look back into the past and a view to the future, *Thromb Haemost* **98** (2007), 49–54.
- [88] A.T. Nurden, Platelets, inflammation and tissue regeneration, *Thromb Haemost* **105** (Suppl 1) (2011), S13–S33.
- [89] S. Offermanns, Activation of platelet function through G protein-coupled receptors, *Circulation Research* **99** (2006), 1293–1304. 10.1161/01.RES.0000251742.71301.16.
- [90] M.A. Packham and M.L. Rand, Historical perspective on ADP-induced platelet activation, *Purinergic Signalling* **7** (2011), 283–292. 10.1007/s11302-011-9227-x.
- [91] J. Papp, P. Kenyeres and K. Toth, Clinical importance of antiplatelet drugs in cardiovascular diseases, *Clinical Hemorheology and Microcirculation* **53** (2013), 81–96. 10.3233/CH-2012-1578.

- [92] J.Y. Park and J.E. Davies, Red blood cell and platelet interactions with titanium implant surfaces, *Clinical Oral Implants Research* **11** (2000), 530–539. 10.1034/j.1600-0501.2000.011006530.x.
- [93] J.Y. Park, C.H. Gemmell and J.E. Davies, Platelet interactions with titanium: Modulation of platelet activity by surface topography, *Biomaterials* **22** (2001), 2671–2682. 10.1016/s0142-9612(01)00009-6.
- [94] C.G. Peters, A.D. Michelson and R. Flaumenhaft, Granule exocytosis is required for platelet spreading: Differential sorting of alpha-granules expressing VAMP-7, *Blood* **120** (2012), 199–206. 10.1182/blood-2011-10-389247.
- [95] S.R. Piersma, H.J. Broxterman, M. Kapci, R.R. de Haas, K. Hoekman, H.M. Verheul and C.R. Jimenez, Proteomics of the TRAP-induced platelet releasate, *J Proteomics* **72** (2009), 91–109.
- [96] G. Pietramaggiore, S.S. Scherer, J.C. Mathews, T. Gennaoui, L. Lancerotto, G. Ragno, C.R. Valeri and D.P. Orgill, Quiescent platelets stimulate angiogenesis and diabetic wound repair, *Journal of Surgical Research* **160** (2008), 169–177. 10.1016/j.jss.2008.09.010.
- [97] W.S. Pietrzak and B.L. Eppley, Platelet rich plasma: Biology and new technology, *J Craniofac Surg* **16** (2005), 1043–1054.
- [98] W.K. Pollock, T.J. Rink and R.F. Irvine, Liberation of [<sup>3</sup>H]arachidonic acid and changes in cytosolic free calcium in fura-2-loaded human platelets stimulated by ionomycin and collagen, *The Biochemical Journal* **235** (1986), 869–877.
- [99] T.M. Quinton, S. Murugappan, S. Kim, J. Jin and S.P. Kunapuli, Different G protein-coupled signaling pathways are involved in alpha granule release from human platelets, *Journal of Thrombosis and Haemostasis: JTH* **2** (2004), 978–984. 10.1111/j.1538-7836.2004.00741.x.
- [100] W.H. Reinhart, Platelets in vascular disease, *Clinical Hemorheology and Microcirculation* **53** (2013), 71–79. 10.3233/CH-2012-1577.
- [101] C.S. Rinder, L.A. Student, J.L. Bonan, H.M. Rinder and B.R. Smith, Aspirin does not inhibit adenosine diphosphate-induced platelet alpha-granule release, *Blood* **82** (1993), 505–512.
- [102] T.J. Rink, Cytosolic calcium in platelet activation, *Experientia* **44** (1988), 97–100.
- [103] T.J. Rink and T.J. Hallam, What turns platelets on, *Trends in Biochemical Sciences* **9** (1984), 215–219. 10.1016/0968-0004(84)90067-7.
- [104] T.J. Rink and S.O. Sage, Calcium signaling in human platelets, *Annu Rev Physiol* **52** (1990), 431–449. 10.1146/annurev.ph.52.030190.002243.
- [105] T.J. Rink, A. Sanchez and T.J. Hallam, Diacylglycerol and phorbol ester stimulate secretion without raising cytoplasmic free calcium in human platelets, *Nature* **305** (1983), 317–319.
- [106] T.J. Rink, S.W. Smith and R.Y. Tsien, Cytoplasmic free Ca<sup>2+</sup> in human platelets: Ca<sup>2+</sup> thresholds and Ca-independent activation for shape-change and secretion, *FEBS letters* **148** (1982), 21–26.
- [107] L. Rivadeneyra, A. Carestia, J. Etulain, R.G. Pozner, C. Fondevila, S. Negrotto and M. Schattner, Regulation of platelet responses triggered by Toll-like receptor 2 and 4 ligands is another non-genomic role of nuclear factor-kappaB, *Thrombosis Research* **133** (2014), 235–243. 10.1016/j.thromres.2013.11.028.
- [108] D. Ronnlund, Y. Yang, H. Blom, G. Auer and J. Widengren, Fluorescence nanoscopy of platelets resolves platelet-state specific storage, release and uptake of proteins, opening up future diagnostic applications, *Advanced Healthcare Materials* **1** (2012), 707–713. 10.1002/adhm.201200172.
- [109] G.J. Roth, A new “kid” on the platelet thrombin receptor “block”: Glycoprotein Ib-IX-V, *Proc Natl Acad Sci U S A* **98** (2001), 1330–1331. 10.1073/pnas.98.4.1330.
- [110] Z.M. Ruggeri, Platelets in atherothrombosis, *Nat Med* **8** (2002), 1227–1234.
- [111] V. Rumjantseva, P.K. Grewal, H.H. Wandall, E.C. Josefsson, A.L. Sorensen, G. Larson, J.D. Marth, J.H. Hartwig and K.M. Hoffmeister, Dual roles for hepatic lectin receptors in the clearance of chilled platelets, *Nature Medicine* **15** (2009), 1273–1280. 10.1038/nm.2030.
- [112] V. Rumjantseva and K.M. Hoffmeister, Novel and unexpected clearance mechanisms for cold platelets, *Transfusion and Apheresis Science: Official Journal of the World Apheresis Association: Official Journal of the European Society for Haemapheresis* **42** (2010), 63–70. 10.1016/j.transci.2009.10.008.
- [113] M. Schattner and G.A. Rabinovich, Galectins: New agonists of platelet activation, *Biol Chem* **394** (2013), 857–863. 10.1515/hsz-2013-0108.
- [114] A. Schedel, S. Thornton, P. Schloss, H. Kluter and P. Bugert, Human platelets express functional alpha7-nicotinic acetylcholine receptors, *Arteriosclerosis, Thrombosis, and Vascular Biology* **31** (2011), 928–934. 10.1161/ATVBAHA.110.218297.
- [115] D.J. Schneider, P.B. Tracy, K.G. Mann and B.E. Sobel, Differential effects of anticoagulants on the activation of platelets *ex vivo*, *Circulation* **96** (1997), 2877–2883.

- [116] S.W. Schneider, S. Nuschele, A. Wixforth, C. Gorzelanny, A. Alexander-Katz, R.R. Netz and M.F. Schneider, Shear-induced unfolding triggers adhesion of von Willebrand factor fibers, *Proc Natl Acad Sci U S A* **104** (2007), 7899–7903. 10.1073/pnas.0608422104.
- [117] A. Schomig, F.J. Neumann, A. Kastrati, H. Schuhlen, R. Blasini, M. Hadamitzky, H. Walter, E.M. Zitzmann-Roth, G. Richardt, E. Alt, C. Schmitt and K. Ulm, A randomized comparison of antiplatelet and anticoagulant therapy after the placement of coronary-artery stents, *The New England Journal of Medicine* **334** (1996), 1084–1089. 10.1056/NEJM199604253341702.
- [118] A. Schomig, N. Sarafoff and M. Seyfarth, Triple antithrombotic management after stent implantation: When and how? *Heart* **95** (2009), 1280–1285. 10.1136/hrt.2008.160457.
- [119] K. Schror, Antiplatelet drugs. A comparative review, *Drugs* **50** (1995), 7–28.
- [120] D.W. Scott, M.O. Vallejo and R.P. Patel, Heterogenic endothelial responses to inflammation: Role for differential N-glycosylation and vascular bed of origin, *Journal of the American Heart Association* **2** (2013), e000263. 10.1161/JAHA.113.000263.
- [121] J.W. Semple, J.E. Italiano Jr. and J. Freedman, Platelets and the immune continuum, *Nat Rev Immunol* **11** (2011), 264–274.
- [122] R. Shirakawa, A. Yoshioka, H. Horiuchi, H. Nishioka, A. Tabuchi and T. Kita, Small GTPase Rab4 regulates Ca<sup>2+</sup>-induced alpha-granule secretion in platelets, *J Biol Chem* **275** (2000), 33844–33849. 10.1074/jbc.M002834200.
- [123] S. Simsekylimaz, E.A. Liehn, C. Militaru and F. Vogt, Progress in interventional cardiology: Challenges for the future, *Thromb Haemost* **113** (2015), 464–472. 10.1160/TH14-07-0599.
- [124] A.L. Sorensen, V. Rumjantseva, S. Nayeb-Hashemi, H. Clausen, J.H. Hartwig, H.H. Wandall and K.M. Hoffmeister, Role of sialic acid for platelet life span: Exposure of beta-galactose results in the rapid clearance of platelets from the circulation by asialoglycoprotein receptor-expressing liver macrophages and hepatocytes, *Blood* **114** (2009), 1645–1654. 10.1182/blood-2009-01-199414.
- [125] T.H. Spaet, The platelet in hemostasis, *Annals of the New York Academy of Sciences* **115** (1964), 31–42.
- [126] T.J. Stalker, E.A. Traxler, J. Wu, K.M. Wannemacher, S.L. Cermignano, R. Voronov, S.L. Diamond and L.F. Brass, Hierarchical organization in the hemostatic response and its relationship to the platelet-signaling network, *Blood* **121**, 1875–1885. 10.1182/blood-2012-09-457739.
- [127] B.Z. Stanger and M.L. Kahn, Platelets and tumor cells: A new form of border control, *Cancer Cell* **24** (2013), 9–11.
- [128] S. Tadokoro, T. Nakazawa, T. Kamae, K. Kiyomizu, H. Kashiwagi, S. Honda, Y. Kanakura and Y. Tomiyama, A potential role for alpha-actinin in inside-out alphaIIb beta3 signaling, *Blood* **117** (2011), 250–258. 10.1182/blood-2009-10-246751.
- [129] J.-C. Tardif, Coronary artery disease in 2010, *European Heart Journal Supplements* **12** (2010), C2–C10. 10.1093/eurheartj/suq014.
- [130] M.L. Taylor, N.L.A. Misso, G.A. Stewart and P.J. Thompson, Differential expression of platelet activation markers Cd62p and Cd63 following stimulation with Paf, arachidonic-acid and collagen, *Platelets* **6** (1995), 394–401. 10.3109/09537109509078478.
- [131] T. Tolmachova, M. Abrink, C.E. Futter, K.S. Authi and M.C. Seabra, Rab27b regulates number and secretion of platelet dense granules, *Proceedings of the National Academy of Sciences of the United States of America* **104** (2007), 5872–5877.
- [132] T. Tsuji and T. Osawa, The carbohydrate moiety of human platelet glycosialin: The structures of the major Asn-linked sugar chains, *Journal of Biochemistry* **101** (1987), 241–249.
- [133] T. Tsuji, S. Tsunehisa, Y. Watanabe, K. Yamamoto, H. Tohyama and T. Osawa, The carbohydrate moiety of human platelet glycosialin, *The Journal of Biological Chemistry* **258** (1983), 6335–6339.
- [134] H. van Nispen tot Pannerden, F. de Haas, W. Geerts, G. Posthuma, S. van Dijk and H.F. Heijnen, The platelet interior revisited: Electron tomography reveals tubular alpha-granule subtypes, *Blood* **116** (2010), 1147–1156. 10.1182/blood-2010-02-268680.
- [135] J.E. Vance and G. Tasseva, Formation and function of phosphatidylserine and phosphatidylethanolamine in mammalian cells, *Biochimica et Biophysica Acta* **1831** (2013), 543–554. 10.1016/j.bbalip.2012.08.016.
- [136] D. Varga-Szabo, A. Braun and B. Nieswandt, Calcium signaling in platelets, *J Thromb Haemost* **7** (2009), 1057–1066.
- [137] D. Varga-Szabo, A. Braun and B. Nieswandt, STIM and Orai in platelet function, *Cell Calcium* **50** (2011), 270–278.
- [138] A. Vieira-de-Abreu, R.A. Campbell, A.S. Weyrich and G.A. Zimmerman, Platelets: Versatile effector cells in hemostasis, inflammation, and the immune continuum, *Semin Immunopathol* **34** (2012), 5–30.
- [139] D.D. Wagner and P.S. Frenette, The vessel wall and its interactions, *Blood* **111** (2008), 5271–5281. 10.1182/blood-2008-01-078204.
- [140] H.H. Wandall, V. Rumjantseva, A.L. Sorensen, S. Patel-Hett, E.C. Josefsson, E.P. Bennett, J.E. Italiano Jr., H. Clausen, J.H. Hartwig and K.M. Hoffmeister, The origin and function of platelet glycosyltransferases, *Blood* **120** (2012), 626–635.

- [141] E.R. Weibel, Fifty years of Weibel-Palade bodies: The discovery and early history of an enigmatic organelle of endothelial cells, *Journal of Thrombosis and Haemostasis: JTH* **10** (2012), 979–84. 10.1111/j.1538-7836.2012.04718.x.
- [142] H.J. Weiss, V.T. Turitto and H.R. Baumgartner, Effect of shear rate on platelet interaction with subendothelium in citrated and native blood. I. Shear rate–dependent decrease of adhesion in von Willebrand’s disease and the Bernard-Soulier syndrome, *The Journal of Laboratory and Clinical Medicine* **92** (1978), 750–764.
- [143] J.G. White and G. Escolar, EDTA-induced changes in platelet structure and function: Adhesion and spreading, *Platelets* **11** (2000), 56–61.
- [144] S.W. Whiteheart, Platelet granules: Surprise packages, *Blood* **118** (2011), 1190–1191.
- [145] D.H. Whitman, R.L. Berry and D.M. Green, Platelet gel: An autologous alternative to fibrin glue with applications in oral and maxillofacial surgery, *J Oral Maxillofac Surg* **55** (1997), 1294–1299.
- [146] P. Wijten, T. van Holten, L.L. Woo, O.B. Bleijerveld, M. Roest, A.J.R. Heck and A. Scholten, High precision platelet releasate definition by quantitative reversed protein profiling–brief report, *Arterioscler Thromb Vasc Biol Dec* **33** (2013), 1635–1638. 10.1161/atvbaha.113.301147.
- [147] O. Yasuda, Y. Takemura, H. Kawamoto and H. Rakugi, Aspirin: Recent developments, *Cellular and Molecular Life Sciences* **65** (2008), 354–358. 10.1007/s00018-007-7449-4.
- [148] M.R. Yeaman, Platelets in defense against bacterial pathogens, *Cellular and Molecular Life Sciences* **67** (2010), 525–544. 10.1007/s00018-009-0210-4.
- [149] A. Yoshioka, H. Horiuchi, R. Shirakawa, H. Nishioka, A. Tabuchi, T. Higashi, A. Yamamoto and T. Kita, Molecular dissection of alpha- and dense-core granule secretion of platelets, *Atherosclerosis Vi* (2001), 403–406.
- [150] A. Zufferey, P. Fontana, J.L. Reny, S. Nolli and J.C. Sanchez, Platelet proteomics, *Mass Spectrometry Reviews* **31** (2012), 331–351. 10.1002/mas.20345.
- [151] R.F. Zwaal, P. Comfurius and E.M. Bevers, Lipid-protein interactions in blood coagulation, *Biochimica et Biophysica Acta* **1376** (1998), 433–453.